J-Psi and the New Physics

G. Preparata, Geneva

(Secretary : High Energy and Particle Physics Division)

The first day of November 1974 saw a remarkable event happening in the world of high energy particle physics. Almost simultaneously, two groups : one led by Sam Ting working at the Brookhaven National Laboratory and the other by Burt Richter at the Stanford Linear Accelerator reported the discovery of a new particle : the J- Ψ (J was the name Ting chose for his discovery, while the Stanford group christened it Ψ). The clarity of their discovery is seen in Figs. 1 and 2.

The finding of the J- Ψ brought finally to fruition two lines of research which were initiated more than a decade ago and are likely to be even more fruitful in the future. Ting's experiment was based on the search for electron-positron pairs produced in high energy proton-beryllium collisions. The first pioneering experiments of such type were carried out in 1965 by Zichichi and his group at CERN. It is only through the application of the technique used to discriminate pions from electrons that Ting could find the true needle in the haystack which the J- Ψ represents in these searches. A similar experiment searching for $\mu^+ \mu^-$ pairs rather than $e^+ e^-$ pairs done by Lederman and his group at BNL had failed to establish the existence of J- Ψ .

The other line of research consists in bringing into collision beams of electrons and positrons and using their energy to produce all sorts of things ; pions, electrons, protons and then in addition the J- Ψ . In the early sixties, the first such machine to be shown to work was built by B. Touschek and his group at Frascati and was called ADA (Anello d'Accumulazione). This was followed by a machine developed by a Stanford-Princeton collaboration

of which Richter was a member. Since then, Orsay and Novosibirsk built higher energy e^+e^- machines, ADA became Adone, DESY got DORIS, and finally it was for SPEAR, a superb example of its breed, to realize the full potentiality of the e^+e^- physics.

With the award of the 1976 Nobel Prize for Physics to Sam Ting and Burt Richter, the J- Ψ has been attributed the highest scientific recognition. It is then an appropriate time to try to assess the meaning and the impact on high energy physics of this wonderful discovery.

The Remarkable J- Ψ

The discovery of a new particle is quite routine in high energy physics ; almost every conference witnesses the birth of new members in the ever growing family of the so called "elementary particles" ; today we know about two hundred of them.

What made the discovery of Ting, Richter and their groups so exciting was its extraordinary lifetime. Until then, in fact, particles as heavy as the J- Ψ (3100 MeV, about 3 proton masses) were known to live for very short time (of the order of 10^{-23} s). The J- Ψ on the other hand, it was immediately clear, lived some thousand times longer than normal. It was as if one found in some remote region of the earth a human race whose life expectancy was not 70 but rather 70 000 years ! Something really unbelievable.

Had the Neutral W-boson Been Found ?

The first reactions of the physics community were of incredulity that the J- Ψ could have anything to do with the dozens of known particles, brought to life by the big accelerators. It had to be something very different ! Could

it be the ever elusive neutral W-boson ? Such a particle along with its charged partners had been postulated to account for the similarities between electromagnetism and weak interactions (responsible among other things for the neutron decay) : a sort of heavy photon.

With not too much effort, one could account for its long lifetime and for its production cross section in both the BNL and SPEAR experiments, but other results from SPEAR did not support the hypothesis. The J- Ψ showed no sign of parity violation, a major feature of the weak interaction. Another blow, probably the fatal one to the W-boson interpretation, was the observation of J- Ψ production in the scattering of high energy photons with beryllium nuclei, observed by W. Y. Lee and his group working at the National Accelerator Laboratory (NAL) in Batavia. The substantial J- Ψ production cross section observed in this experiment could in no way be accounted for by weak interactions. On the other hand, if the J- Ψ was assumed to be a vector meson, similar in nature to the well known Φ -meson, (discovered in the early sixties), reasonable agreement with the experimental findings could be obtained.

Charm : the New Quantum Number

Once the W-hypothesis was out of the way, the odds increased for another hypothesis which was pursued very vigorously by Shelly Glashow and his Harvard Group. They had been stressing since its discovery that the J- Ψ was indeed a hadron (the generic name for particles which interact strongly, like the π -meson and the proton) and its remarkable lifetime was related to the existence for hadronic matter of a new quantum number be-

Left - Samuel Ting

Nobel Prize Winners in 1976

Right - Burton Richter

Fig. 1 The J/ψ signal detected by Ting and his group at BNL. The J/ψ appears as a quite striking peak in the distribution of the total mass of the electron-positron pairs.

sides Isotopic spin and Strangeness: **Charm**. Actually this new quantum number had appeared on and off in the scientific literature since the confirmation of the Gell-Mann Ne'eman SU (3) scheme, by the discovery of that very peculiar member of the zoo of elementary particles which is the Ω^- . From 1963 on, physicists had been wondering whether that quality of hadrons called "Strangeness" was indeed unique; could not there be some other kind of super-strangeness or Charm? The works of Hara, Bjorken and Glashow, and Amati, Bacry, Nuyts and Prentki were the first attempts to introduce Charm into the hadronic world, and the idea rested in peace until in 1970 it was resurrected by Glashow, Iliopoulos and Maiani in the context of a model for the weak interactions of the hadrons (GIM model).

The revived interest in the gauge theory of weak interactions proposed by Salam and Weinberg (SW model) which was sparked off by the 't Hooft proof (1971) of its renormalizability (i.e.

Fig. 3 The naive quark model description of two well-known hadrons: the π^+ (left) and the proton (right).

the possibility of removing troublesome infinities in perturbative calculations) and the discovery of neutral currents by the "Gargamelle" group working at CERN, involved also the GIM model as a possible extension of the SW-model to the hadronic weak interactions and gave further motivation for the search for such a quantum number as emphasized particularly by Gaillard, Lee and Rosner (1974). Thus in 1974 everybody was ready to welcome the discovery of Charm. But then, why did the J/ψ cause so much confusion? The answer is quite simple: nobody was prepared to describe as hadronic, an object which lived some thousand times longer than normal. Some lengthening of its lifetime was indeed expected, by following through the analogy with the Φ -meson, but nothing of the size that was observed.

However the elimination of alternative hypotheses (besides the W-boson, another candidate soon fallen in disgrace had been the theory of Colour proposed in 1965 by Han and Nambu) corroborated the suspicion that with the J/ψ , a new quantum number had appeared in the world of hadrons. It was now a matter of further experimental work to confirm or reject this suspicion.

The New vs. the Old Physics

We cannot understand what physics vistas, the "New Physics", i.e. the J/ψ discovery has been opening up if we do not try to set the stage of the "Old Physics" that high energy physicists have been investigating over the past 30 years, since the momentous discovery of the π -meson. It must be said that in spite of the enormous amount of information available today on the most diverse aspects of hadrons and their interactions, the "Old Physics" is far from being a closed chapter of natural science. We do not yet know the laws obeyed by this peculiar kind of matter, nor can we calculate with any confidence some of its simplest properties. We do possess, however, a set of rules which can relate and organize an impressive body of experimental information in a simple and amazingly successful way: The Quark Model.

According to the quark model the hadronic particles are all built up from basic constituents, the quarks, which carry the fundamental quantum numbers of the world of hadrons. Before the J/ψ , the observations were consistent with the existence of three spin- $1/2$ objects: p , n and λ (also called u , d , s), whose (Charge, Strangeness) assignments are $(2/3, 0)$, $(-1/3, 0)$ and $(-1/3, -1)$ respectively. One manu-

Fig. 2 This is how the SPEAR group saw the J/ψ . In (a) the e^+e^- annihilation cross-section increases by a factor hundred in passing through the resonance at 3100 MeV; (b) and (c) show the J/ψ signal in the e^+e^- and $\mu^+\mu^-$ channels respectively.

factures hadrons by combining either a quark and an antiquark (Meson) or three quarks (Baryons); much in the same way as the positronium is composed of an electron-positron pair and the nucleus of ^3He is made out of three nucleons. The picture above is supported to a high degree by our knowledge of the spectrum of hadrons: all known particles without exception can be described as quantum mechanical states of such physical systems. This is very nice, but strangely enough, the more the quark model unravels about hadrons, the further it leads us towards an inextricable paradox. For if the hadrons are composite, under appropriate observational conditions their constituents should reveal themselves; this is what happens for any atom or nucleus. Nobody, on the other hand, has yet been able to isolate quarks. An easy way out, which comes to mind at once, is that quarks have such a high mass that they cannot be produced as free objects until we dispose of the appropriate energy (and this in practice may never happen!). Even

discounting adverse geo-chemical evidence, the high quark mass appears like a stratagem which poses more puzzles than it solves. For in a situation of enormous binding energy such as is required by the previous "explanation", it is unbelievable that the simple dynamics implied by the quark model can manage to survive. The monstrously strong forces acting between them, will produce incessantly quark-antiquark pairs and with them, physical systems whose structure is infinitely more complicated than the one conceived by the simple quark model.

In spite of the quark paradox, whose resolution will most likely require radically new ideas, the successes of the quark model have generated a beneficial degree of confidence that the ideas developed for the "old observed quarks" (as p , n , λ have optimistically been called) can be extended to describe the Charm quantum number, by simply introducing a new **charmed** quark, whose charge is $2/3$, its isospin zero, and its strangeness zero. "The New Physics" deals with the states where at least one of the quarks is a c -quark (or anti c -quark) much in the same way as the "Old Physics" deals with the hadrons entirely made out of p , n , and λ quarks (antiquarks).

The New Physics

Equipped with the theoretical apparatus of the quark model, the New Physics could set its main experimental goals which are being achieved with considerable success. In the quark picture, the $J-\Psi$ is a vector meson built up by a $c\bar{c}$ -pair in S -wave, carrying a total quark-spin $|S_q|=1$. (It is in spectroscopic notation a 3S_1 state). Its ridiculously small width (long lifetime) was, and still remains, a big puzzle; it conceivably involves some dynamical aspect which the quark model, in its admitted coarseness, cannot handle realistically. As a member of the $c\bar{c}$ family, the $J-\Psi$ is expected to have brothers with well defined characteristics like the 3P -states, some of which for old particles as the Φ have been observed [i.e. the $S^*(0^{++})$ and the $f^*(2^{++})$]. If the dynamics is not too crazy, these states should lie in a mass range between the $J-\Psi$ and its first radial recurrence, the Ψ' , which was observed at a mass $M=3965$ MeV at SPEAR, soon after the $J-\Psi$.

A careful and difficult analysis of γ -spectra carried out both at SPEAR and at DESY (Hamburg), produced in the Summer of 1975, direct evidence of such states (called Ξ 's) in the mass range 3400-3500 MeV: another very

good point in favour of the new quark interpretation. At the Stanford Conference (August '75), DESY reported evidence of another state at around 2800 MeV, which could be interpreted as the 3S_0 partner of the $J-\Psi$, and was therefore called η_c (recalling the well known η particle of "old physics").

Furthermore refined studies of the total e^+e^- annihilation cross section carried out at SPEAR, revealed very suggestive bumps in the energy range between 4000 and 4500 MeV. They could be interpreted as higher vector members of the $J-\Psi$ family, their much bigger width being due to a different decay mechanism which became available if the charmed particle threshold lay below 4000 MeV. This was a very important hint of where to look for the so far elusive charmed particles, for until then, the evidence for Charm had been either indirect or quite fuzzy, coming from the observation of the $J-\Psi$ and its family and from some strange neutrino events (we shall return to them below).

We must note that the Charm of the $J-\Psi$ is only "hidden", its total charm in fact is zero, the Charm of its c -quark being cancelled by that of its anti c -quark. But in a quark model, truly charmed states, corresponding to configurations like $c\bar{p}$ or $c\bar{n}$ (D) and $c\bar{\lambda}$ (F), must exist and their mass lie somewhere near 2000 MeV. These states, due to Charm-conservation in strong and electromagnetic interactions, can decay only through weak interactions with lifetimes which can be estimated around 10^{-13} - 10^{-12} s.

Thus, if these ideas are correct, under the bumps seen at SPEAR above 4000 MeV, there had to lie plenty of charmed particles, waiting only for experimental ingenuity to sort them out. In the Spring '76 the first clear evidence for D -type states was presented in the form of a striking ($K\pi$) peak with a mass of 1850 MeV. A $K\pi\pi$ state was also reported at about the same mass, evoking the old puzzles that the idea of Strangeness many years ago helped to resolve. Thus charmed particles had been positively established and the hypothesis of a new quark had met with yet another crucial corroboration.

New Physics and the Neutrino

As mentioned in passing above, not all the action of the New Physics took place in strong and electromagnetic forms. Soon after the $J-\Psi$ had been discovered, a Brookhaven group found in a bubble chamber exposed to a beam of neutrinos, one example of the reaction

$$\nu_\mu p \rightarrow \mu^- \Lambda^0 \pi^+ \pi^+ \pi^-$$

This process contradicts the firmly established $\Delta Q = \Delta S$ rule of weak interactions, according to which the variation of the hadrons' Charge between the initial and the final state must equal the variation of their Strangeness. The BNL group suggested that this anomaly was only apparent, and that what one was really observing was the decay of a neutrino-produced charmed baryon (a pnc configuration) whose mass was determined to be about 2450 MeV: the expected ball park for such an object!

Other possible evidence for Charm neutrino production was supplied by the Gargamelle group at CERN, who observed a few events of the type:

$$\nu_\mu N \rightarrow \mu^- e^+ + \text{hadrons},$$

which could be interpreted as the production and the subsequent leptonic decay of a charmed baryon. As for very high energy neutrinos, such as those manufactured by the NAL machine, they too seem to produce quite easily charmed particles, as indicated by the relatively numerous dimuon events, i.e.

$$\nu_\mu \text{Nucleus} \rightarrow \mu^+ \mu^- + \text{hadrons}$$

observed by both the Harvard-Penn-Wisconsin and the Caltech groups working at Batavia. Actually high energy neutrino experiments may reveal themselves extraordinarily rich mines of new objects; several "anomalies" are being reported from NAL whose explanation might require yet another extension of the hadronic quantum numbers as well as of the structure of the weak interactions. This is why today high energy physicists are looking very keenly at neutrino interactions; very soon the 400 GeV SPS accelerator at CERN will lead the world effort in this fascinating field.

Lessons and Expectations

After this necessarily sketchy outline of the $J-\Psi$ discovery and the emergence of the New Physics, let me conclude with a few reflections.

Looking back over the two years that have elapsed since the $J-\Psi$ was discovered, we can draw a number of lessons. The first and most important is that the quark model has emerged as a truly predictive theoretical tool, embodying a great deal of truth about hadrons and their interactions. The quark ideas had to stand most stringent tests like the existence of the P -States, the charmed particles and some intriguing aspects of their weak decays (which cannot be discussed here), and have passed them so handsomely that it would be astonishing if all this were a mere accident. Never before have we been so confident that the quark model is here to stay, and

the New Physics has been crucial in strengthening our confidence. But can the New Physics help us to solve the tantalizing difficulties of the paradoxical quark behaviour?

Many physicists have expressed strong hopes that the spectrum and transition rates of the $J-\Psi$ family, the "Charmonium" system as it has been called (paraphrasing the "positronium") can be to the finding of the correct quark theory, what the hydrogen atom has been to the formulation of Quantum Mechanics. Another lesson we have learned is that, through the quark concept, the weak interactions exhibit a remarkable symmetry between leptons and hadrons. It will be very interesting to see whether this symmetry survives the possible discovery of new types of leptons and/or hadrons (some peculiar events found at SPEAR indicate that a heavy lepton has already been found).

It is really within high energy weak interactions, with their quark-lepton symmetry, that matter could one day show truly universal features, thus leading to a unification of all forms of interactions. Such grandiose schemes are presently being conceived, a legitimate offspring of the New Physics.

Who knows; is the $J-\Psi$ the first link of a chain leading to the fulfilment of the ancient aspiration to cosmic unity?

Instituut voor Kernfysisch Onderzoek

I KO

Applications are invited for the position of :

Groupleader

in Pion-Muon Nuclear Physics

at the Institute for Nuclear Physics Research (IKO), Amsterdam.

The Institute is in the course of constructing a high duty cycle 500 MeV linear electron accelerator and several associated facilities.

Qualifications	— experience with large and complex instrumentation — relevant research experience at senior level — team leadership and management experience — sufficient scientific background to guide research in front-line of physics.
Responsibilities	— construction of a magnetic muon channel — planning of experimental equipment for π/μ facility — guidance of both the research and the construction teams.

A salary will be offered in the range between Dfl. 5000,— and Dfl. 6300,— per month, depending on age and experience.

Enquiries should be addressed to Dr. P.F.A. Goudsmit, Institute for Nuclear Physics Research, P.O. Box 4395, Amsterdam, The Netherlands (telephone 020-930951).

Applications, including a curriculum vitae, list of publications and a least two references should be made before March 15, 1977.

Physics in Hungary

N. Kroó, Budapest

(Hungarian Academy of Sciences)

The study of physical problems has always played an important role in the scientific life of Hungary. Outstanding researchers, Loránd Eötvös being a very good example, achieved remarkable results, though mostly in difficult circumstances. Unfortunately, many talented Hungarian physicists, such as Kúrmán, Hevessy, Szilárd, Wigner, and others found it necessary to continue their scientific careers abroad.

Although the roots of the present scientific activities can be traced back to researches carried out in the 1920s and 1930s, really rapid development started only after the Second World War. Physics research is carried on in Hungary, as in many other countries, in three types of laboratory : university departments, industrial research institutes and in institutes of the Hungarian Academy of Sciences (HAS). The most

spectacular change has been the evolution of the Academy institutes, as is apparent from the figures listed below:

	1950	1960	1970
Number of physicists in HAS institutes	6	186	310
Relative allocation per physicist to physics research	1	7	8

While the Academy plays an important role in directing research, the Eötvös Loránd Physical Society is an important social organization enabling physicists from educational centres, research institutes and from industry to exchange ideas in the conferences, schools and in the club-like lecture meetings organized by the Society which is the representative association of Hungarian physicists.

Activities in Physics

Studies in modern theoretical physics started in Hungary in the early thirties. Quantum theory, statistical mechanics and the theory of relativity were the main fields of research. A series of regular high-level seminars with participants from home and abroad helped to promote an interest in these subjects.

After the Second World War the initial task was to lay the foundation of future development by emphasizing the importance of an education in theoretical physics. Simultaneously, a high level school in quantum field theory, that still functions, was established.

The statistical theory of atoms, the study of which was initiated in this early period, is still a subject pursued