

nedm.web.psi.ch  
www.edm.ethz.ch

© ILL

# SEARCHES FOR NEW PHYSICS WITH FREE NEUTRONS AND RADIOACTIVE ATOMIC NUCLEI

■ N. Severijns – Instituut voor Kern- en Stralingsfysica, Katholieke Universiteit Leuven, Belgium – DOI: <https://doi.org/10.1051/epn/2021405>

The Standard Model of Particle Physics is very successful but does not explain several experimental observations. Extensions of it, invoking new particles or phenomena, could overcome this. Experiments in different energy domains allow testing these extensions and searching for new particles. Here focus is on low-energy experiments with neutrons and radioactive nuclei.

## The Standard Model of Particle Physics and beyond

The Standard Model of Particle Physics (SM) successfully describes the fundamental particles (quarks and leptons), and their interactions via the electromagnetic and the weak and strong nuclear forces by exchanging messenger particle ‘gauge bosons’ (the photon for

electromagnetism). However, it does not explain a number of observed phenomena. Examples are the breaking of parity symmetry by the weak force (meaning that of a weak decay process and its three-dimensional mirror image only one effectively occurs in Nature), the precise nature of this force, and the difference between the amount of matter and antimatter in the Universe.

▲ nEDM inner view vacuum chamber. © ILL

Extensions of the SM try explaining these phenomena by adding specific features or new gauge bosons. Experiments try to confirm or falsify these extensions, indicating the direction for future work. Best known are searches for new particles at high-energy colliders. A good example is the discovery of the Higgs-boson (related to the generation of the mass of all particles), at the Large Hadron Collider at CERN (Geneva). Here we focus on experiments at much lower energies, with free neutrons and radioactive atomic nuclei. Recent reviews of this field are found in *e.g.* [1-2].

### The matter-antimatter mystery and electric dipole moments

A currently very active field of research is trying to shed light on the small difference between the amount of matter and antimatter that must have occurred in the very early universe. When all antimatter had disappeared in ‘collisions’ with matter particles, thereby generating the cosmic background radiation we observe today, a small excess of matter (about one particle in a billion) evolved to form the universe we observe today, and is the cause for us being here. Andrei Sakharov showed that such a difference can be explained by three conditions, one of these being a violation of the so-called CP symmetry. This is the combination of charge symmetry (C) and parity symmetry (P), which requires replacing in a physical process all particles by their antiparticles (C) and taking the (three-dimensional) mirror image of the process (P). If the resulting process is also observed in Nature, CP symmetry is said to hold, if not it is violated. Violation of CP-symmetry is observed in a certain class of particles (called  $K^0$ ,  $B^0$  and  $D^0$  mesons), but this is too small to explain the observed matter-antimatter difference. Physicists therefore search for other, larger signals of CP violation.

A very popular observable violating CP symmetry is a permanent electric dipole moment (EDM),  $d$ , of a particle. First experiments started with the neutron in the 1950’s (Fig. 1), and sensitivity has increased by about six orders of magnitude. The current best value was obtained in an experiment at the Paul Scherrer Institute (Switzerland). The result,  $d_n = (0.0 \pm 1.1_{\text{stat}} \pm 0.2_{\text{sys}}) \times 10^{-26}$  e.cm [3], is consistent with zero. The high sensitivity is illustrated by the fact that if one would enlarge the neutron to the size of the Earth, the difference between the time-averaged position of positive and negative charges inside it should be less than several micrometers.

Permanent EDMs are also searched for in other particles, in atoms and in molecules [4]. Indeed, as many extensions of the SM include different CP-violating sources, these can only be distinguished when a variety of systems is studied. The highest-precision EDM results to date were obtained with  $^{199}\text{Hg}$  and  $\text{ThO}$  molecules. Recently, interest has grown in octupole deformed



**Experiments in beta decay have led Pauli to suggest the existence of the neutrino and uncovered several fundamental properties of the weak force, such as the violation of parity symmetry and the overall structure of the weak interaction.”**

nuclei as this was shown to enhance a possible EDM. As such isotopes are found in elements like Rn, Ra, Th, and Pa, with only radioactive isotopes, experiments require radioactive beam facilities, *e.g.* ISOLDE at CERN, Geneva or TRIUMF at Vancouver.

### Studying the weak nuclear force in neutron and nuclear beta decay

The most elementary beta-decay process is that of the neutron, changing it into a proton via the creation of a W gauge boson (a weak force messenger particle), and causing emission of an electron ●●●





Fig. 1: a non-zero permanent EDM,  $d$ , for the neutron would mean that the time-averaged position of positive and negative charges inside the (neutral) neutron would not coincide. This would cause a precession of the neutron spin when placed in an electric field,  $\mathbf{E}$ . As the neutron has a magnetic moment,  $\mu$ , which does not violate CP symmetry and causes a spin precession in a magnetic field,  $\mathbf{B}$ , a non-zero EDM would cause a tiny difference in the neutron precession frequency when placed in alternately parallel (left) and anti-parallel (right) magnetic and electric fields.

••• and an electron-antineutrino ( $\beta^-$  decay). This occurs for a free neutron and for neutrons in radioactive nuclei with an excess of neutrons over protons. In radioactive nuclei with a proton excess, the inverse process ( $\beta^+$  decay) can occur, changing a proton into a neutron with emission of a positron and electron-neutrino. Experiments in beta decay have led Pauli to suggest the existence of the neutrino and uncovered several fundamental properties of the weak force, such as the violation of parity symmetry and the overall structure of the weak interaction, *i.e.* the type of messenger particles that relate to it.

As the neutron (lifetime  $\sim 15$  minutes) is a very simple system consisting of three quarks, it is an ideal probe for studying the weak force with few disturbing effects. In the beta decay of radioactive nuclei (with half-lives from a fraction of a second to billions of years), effects of the nuclear medium (*e.g.* virtual pion exchange) have to be included. However, the several thousand radioactive nuclei have very different properties, so that one can select the ones best suited for a given experiment. Important classes are those with an equal number of protons and neutrons, or a difference of just one (mirror nuclei).

One type of experiments determines the  $ft$ -value of a beta transition, which depends on its intensity and decay energy, and the half-life of the decaying state, corrected for percent-level effects. For so-called pure Fermi transitions this has provided a precise value for the *up-down* quark-mixing matrix element [4]. Similar measurements in neutron decay and for mirror nuclei are not yet competitive but are catching up [2]. The *up-down* matrix element is important for testing unitarity of the quark-mixing matrix, allowing to search for beyond-SM physics such as a fourth generation of quarks or new gauge bosons.

Fig. 2: result of the fit of nuclear and neutron decay data sensitive to scalar and tensor type weak interaction couplings ( $C_S$  and  $C_T$ ) involving right-handed neutrinos, relative to the SM vector and axial-vector coupling constants  $C_V$  and  $C_A$ . The contours of constant  $\chi^2$  correspond to iso- $\chi^2$  levels with values  $\Delta\chi^2 = 1, 4,$  and  $9$ . Based on these limits, scalar and tensor type weak interactions with right-handed neutrinos could in principle occur in up to about 10% of the beta decays. From Ref. [1].



So-called correlation measurements observe the relative emission angle between the spins and momentum vectors of the initial neutron/radioactive nucleus and its decay particles, *i.e.* the electron or positron and the recoiling nucleus (the (anti-)neutrino from beta decay cannot be detected). Almost always the energies of the decay particles are also measured. One can further choose to polarize the spin of the neutron/nucleus in the direction of an applied magnetic field. The forward-backward asymmetry in the electron emission with respect to the polarized neutron spin provides the relative strength of the two types of weak force observed in Nature, the ratio  $g_A/g_V$  [1]. Further, precise tests of the parity or time reversal symmetries [5] can be performed, the latter testing whether a physical process and the ‘in time backward running’ process occur with the same probability. One can, finally, also search for possible new components of the weak force driven by as yet unobserved messenger bosons. The latter would induce a small change in the values of experimental observables with respect to their SM value (*e.g.* Fig. 2). Information obtained is complementary to that from colliders [1, 2].

Of course, instrumental (systematic) effects always have to be carefully considered. In this respect, the possibility to perform extensive simulations of the behaviour of the experimental apparatuses, is a big asset. Together with advances in detection technology and analysis tools, this has allowed reaching precisions at the per mil level. As this requires taking in account also small contributions to the value of observables induced by the strong force (*e.g.* [1]), new challenges and new research possibilities now open up for this field. ■

## About the Author



**Nathal Severijns** is full professor of physics at KU Leuven University with a track record in low-energy weak interaction experiments with spin-polarized and unpolarized neutrons and exotic nuclei. He has written several review articles covering this research field.

## References

- [1] M. González-Alonso, O. Naviliat-Cuncic and N. Severijns, *Prog. Part. Nucl. Phys.* **104**, 165 (2019).
- [2] A. Falkowski, M. González-Alonso, and O. Naviliat-Cuncic, *JHEP* **04**, 126 (2021).
- [3] C. Abel *et al.*, *Phys. Rev. Lett.* **124**, 081803 (2020).
- [4] J.C. Hardy and I.S. Towner, *Phys. Rev. C* **102**, 045501 (2020).
- [5] T. E. Chupp, P. Fierlinger, M. J. Ramsey-Musolf, and J. T. Singh, *Rev. Modern Phys.* **91**, 015001 (2019).

# COMPANY DIRECTORY

Highlight your expertise. Get your company listed in europhysicsnews company directory  
For further information please contact [bernadette.dufour@edpsciences.org](mailto:bernadette.dufour@edpsciences.org)

## BARTINGTON INSTRUMENTS

[www.bartington.com](http://www.bartington.com)

Bartington Instruments designs and manufactures high precision fluxgate magnetometers, Helmholtz coil systems, gradiometers, magnetic susceptibility instruments and associated data processing equipment, for the precise generation and measurement of magnetic fields around physics experiments, and for calibration of magnetometers in consumer electronics.



## BLUEFORS OY

[Bluefors.com](http://Bluefors.com)

Bluefors manufactures cryogenic measurement systems and has a strong focus in the field of quantum technology. Our reliable and easy-to-operate systems can be customized to meet the special requirements of each experiment allowing the customer to get in direct contact with the scientists and engineers who design their system.



## MB SCIENTIFIC AB

[www.mbscientific.se](http://www.mbscientific.se)

MB Scientific AB is a Swedish company which develops and produces state of the art instruments for the photoelectron spectroscopy experiments. Our photoelectron energy analyser MBS A-1 gives you the opportunity to do world leading research together with MBS VUV photon sources, MBS L-1 and T-1, which produce the brightest and narrowest lines existing to be used for this type of experiments.



## MCPHERSON

[www.mcphersoninc.com](http://www.mcphersoninc.com)

McPherson designs and manufactures scanning monochromators, flat-field imaging spectrographs, and vacuum monochromators and measurement systems for reflectance, transmittance, and absorbance testing. Its spectrometers and systems are built for soft x-ray, vacuum-ultraviolet, and UV/Vis and Infrared wavelengths. Applications range from lasers and lithography, solar, and energy to analytical life science and more.



## OXFORD INSTRUMENTS

[nanoscience.oxinst.com](http://nanoscience.oxinst.com)

Oxford Instruments NanoScience provides market-leading cryogenic systems that enable quantum technologies, new materials and device development in the physical sciences. To learn more about Oxford Instruments NanoScience's low temperature systems for quantum computing applications, please visit our website.



## PFEIFFER VACUUM

[www.pfeiffer-vacuum.com/en/](http://www.pfeiffer-vacuum.com/en/)

Pfeiffer Vacuum stands for innovative and custom vacuum solutions worldwide, technological perfection, competent advice and reliable service. With the invention of the turbopump, the company paved the way for further development within the vacuum industry. Pfeiffer Vacuum offers a complete product portfolio: backing pumps, leak detectors, measurement and analysis devices, components as well as vacuum chambers and systems.



## THYRACONT

[thyracont-vacuum.com/en](http://thyracont-vacuum.com/en)

Thyracont develops vacuum gauges and supplies innovative devices for leading manufacturers of vacuum pumps and process plants. As specialists in vacuum metrology, their portfolio includes vacuum sensors, transducers, meters, switches, controllers as well as accessories and components. They likewise offer a wide range of measuring principles in various combinations. Thyracont is involved today in your technological need of tomorrow.



## TOPTICA PHOTONICS

[www.toptica.com](http://www.toptica.com)

TOPTICA Photonics, founded in 1998 near Munich (Germany), develops and manufactures high-end laser systems for scientific and industrial applications. The portfolio includes diode lasers, ultrafast fiber lasers, terahertz systems and frequency combs. OEM customers, scientists, and over a dozen Nobel laureates all acknowledge the world-class exceptional specifications of TOPTICA's lasers, as well as their reliability and longevity.



## ZURICH INSTRUMENTS

[www.zhinst.com](http://www.zhinst.com)

Zurich Instruments is a technology leader developing and selling advanced test & measurement instruments for dynamic signal analysis. These devices are used in many fields of application by high-technology research laboratories and industrial development sites. Zurich Instruments' vision is to revolutionize instrumentation in the high-frequency and ultra-high-frequency range by incorporating the latest analog and digital technology into powerful measurement systems.

